

# ABATACEPT (ORENCIA®) PRESCRIBER ORDER FORM

Patient Name: \_\_\_\_\_ Date of Birth: \_\_\_\_\_ Gender: \_\_\_\_\_

Address: \_\_\_\_\_

Phone: \_\_\_\_\_ Height: \_\_\_\_\_  inches  cm Weight: \_\_\_\_\_  lbs  kg

## Clinical Information

Primary Diagnosis Description: \_\_\_\_\_ ICD-10 Code: \_\_\_\_\_

Allergies:  NKDA OR (List): \_\_\_\_\_

Is this the first dose?  Yes – date of first dose: \_\_\_\_\_ Hepatitis B Status: \_\_\_\_\_ Titer Date: \_\_\_\_\_  
 No – date of last dose: \_\_\_\_\_  Positive  Negative

TB Status:  PPD (negative) – date: \_\_\_\_\_  Active TB  
 Last chest x-ray – date: \_\_\_\_\_  Unknown  
 Past positive TB infection, course taken: \_\_\_\_\_  
 QuantiFERON or T Spot Assay result and date: \_\_\_\_\_

## Abatacept (Orencia®) Prescription

### Abatacept (Orencia®) INTRAVENOUS refill as directed x 1 year

Initial Dose:  Infuse 500 mg IV over 30 minutes on Weeks 0, 2, and 4 (patient weight < 60 kg)  
 Infuse 750 mg IV over 30 minutes on Weeks 0, 2, and 4 (patient weight 60 to 100 kg)  
 Infuse 1000 mg IV over 30 minutes on Weeks 0, 2, and 4 (patient weight > 100 kg)  
 Other: \_\_\_\_\_

Maintenance Dose:  Infuse 500 mg IV over 30 minutes every 4 weeks (patient weight < 60 kg)  
 Infuse 750 mg IV over 30 minutes every 4 weeks (patient weight 60 to 100 kg)  
 Infuse 1000 mg IV over 30 minutes every 4 weeks (patient weight > 100 kg)  
 Other: \_\_\_\_\_

### Abatacept (Orencia®) SUBCUTANEOUS refill as directed x 1 year

Optional loading dose of \_\_\_\_\_ mg IV administered once  
 125mg subcutaneously once weekly (within a day of IV infusion if infusion given)  
 Other: \_\_\_\_\_

## Ancillary Orders

### Anaphylaxis Kit

Dosage:  Epinephrine 0.3 mg (> 30 kg), 0.15 mg (15 to 30 kg), or 0.01 mg/kg (< 15 kg) SQ or IM x 1; repeat x 1 in 5 to 15 min PRN.  
 Diphenhydramine 25 mg (> 30 kg) or 1.25 mg/kg (≤ 30 kg) IV or IM; repeat x 1 in 15 min PRN no improvement.  
 0.9% Sodium Chloride 500 mL (> 30 kg) or 250 mL (≤ 30 kg) IV at KVO rate PRN anaphylaxis.

Skilled Nursing to establish peripheral IV access as needed to manage anaphylaxis

### Medication Orders

Acetaminophen 650 mg PO 30 min before infusion, may repeat every 3 to 4 hours as needed for headache or mild discomfort. Patient may decline.  
 Diphenhydramine 25 mg PO 30 min before infusion, may repeat every 4 to 6 hours as needed for mild to moderate allergic reactions. Patient may decline.  
 Methylprednisolone sodium succinate 40 mg IV push 20 minutes prior to infusion.  
 Other: \_\_\_\_\_

### IV Flush Orders

- Peripheral: 0.9% Sodium Chloride 2 to 3 mL pre-/post-use.
- Implanted Port: 0.9% Sodium Chloride 5 to 10 mL pre-/post-use and 10 to 20 mL pre-/post-lab draw. Heparin (100 unit/mL) 3 to 5 mL post-use. For maintenance, heparin (100 unit/mL) 3 to 5 mL every 24 hr if accessed or weekly to monthly if not accessed.

### Lab Orders

No labs ordered at this time.  
 Other: \_\_\_\_\_

Skilled nurse to assess and administer and/or teach self-administration, where appropriate, via access device as indicated above. Nurse will provide ongoing support as needed. Refill above ancillary orders as directed x 1 year.

If patient is seen within a provider led infusion clinic, Option Care Health's infusion reaction management policy, skilled nursing plan of treatment, and IV flush administration will be followed per provider oversight. No individual anaphylaxis kit will be dispensed.

*I certify that the use of the indicated treatment is medically necessary, and I will be supervising the patient's treatment.*

Prescriber Signature: \_\_\_\_\_ Date: \_\_\_\_\_

## Prescriber Information

Prescriber Name: \_\_\_\_\_ Phone: \_\_\_\_\_ Fax: \_\_\_\_\_

Address: \_\_\_\_\_ NPI: \_\_\_\_\_

City, State: \_\_\_\_\_ Zip: \_\_\_\_\_ Office Contact: \_\_\_\_\_

Fax completed form, insurance information, and clinical documentation to: 713-983-4647

**CONFIDENTIAL HEALTH INFORMATION:** Healthcare information is personal information related to a person's healthcare. It is being faxed to you after appropriate authorization or under circumstances that do not require authorization. You are obligated to maintain it in a safe, secure, and confidential manner. Re-disclosure of this information is prohibited unless permitted by law or appropriate customer/patient authorization is obtained. Unauthorized re-disclosure or failure to maintain confidentiality could subject you to penalties described in federal and state laws. **IMPORTANT WARNING:** This message is intended for the use of the person or entity to whom it is addressed and may contain information that is privileged and confidential, the disclosure of which is governed by applicable law. If the reader of this message is not the intended recipient, or the employee or agent responsible for delivering it to the intended recipient, you are hereby notified that any dissemination, distribution or copying of this information is STRICTLY PROHIBITED. If you have received this message in error, please notify us immediately. Brand names are the property of their respective owners.