

# VEDOLIZUMAB (ENTYVIO®) PRESCRIBER ORDER FORM

Patient Name:		Date of Birth:	Gender:	
Address:				
Phone:	Height:	<input type="checkbox"/> inches <input type="checkbox"/> cm	Weight:	<input type="checkbox"/> lbs <input type="checkbox"/> kg

## Clinical Information

Primary Diagnosis Description:	ICD-10 Code:
Is this the first dose? <input type="checkbox"/> YES – Date of first dose: _____ <input type="checkbox"/> NO – Date of last dose: _____	

## Vedolizumab (Entyvio®) Prescription

Entyvio® (Vedolizumab) refill as directed x 1 year.

Please select one of the options below:

	<input type="checkbox"/> Option 1 IV Regimen:	<input type="checkbox"/> Option 2 IV to SUBQ Regimen:	<input type="checkbox"/> Option 3 IV to SUBQ Regimen:
<b>Induction</b>	Infuse 300 mg IV over at least 30 minutes on Weeks <b>0, 2, 6</b>	Infuse Entyvio 300mg IV over at least 30 minutes at weeks <b>0, 2, 6</b>	Infuse Entyvio 300mg IV over at least 30 minutes at weeks <b>0, 2</b>
<b>Maintenance</b>	Infuse 300 mg IV over at least 30 minutes every 8 weeks thereafter	Inject Entyvio 108mg SUBQ at week 14, then every 2 weeks thereafter	Inject Entyvio 108mg SUBQ at week 6, then every 2 weeks thereafter

Other: \_\_\_\_\_

For IV doses, quantity sufficient of Entyvio® 300mg vials will be dispensed to the patient to complete the prescribed treatment plan. In addition, after each infusion the IV tubing will be flushed with NS 30ml using a 50ml bag.

For SUBQ doses, quantity sufficient of Entyvio® 108mg Prefilled Pens will be dispensed to fulfill prescribed treatment plan.

## Ancillary Orders

### Anaphylaxis Kit

If this is a 1<sup>st</sup> infusion dose, would you like Option Care Health to provide an anaphylaxis kit with the 1<sup>st</sup> dose?

Yes  No

- Dosage:
- Epinephrine 0.3 mg (> 30 kg), 0.15 mg (15 to 30 kg), or 0.01 mg/kg (< 15 kg) SUBQ or IM x 1; repeat x 1 in 5 to 15 min PRN.
  - Diphenhydramine 25 mg (> 30 kg) or 1.25 mg/kg (≤ 30 kg – 25mg max dose) IV or IM; repeat x 1 in 15 min PRN no improvement.
  - 0.9% Sodium Chloride 500 mL (> 30 kg) or 250 mL (≤ 30 kg) IV at KVO rate PRN anaphylaxis.

Skilled Nursing to establish peripheral IV access as needed to manage anaphylaxis.

### Medication Orders

- Acetaminophen 650 mg PO 30 min before infusion, may repeat every 3 to 4 hours as needed for fever or mild discomfort. Patient may decline.
- Diphenhydramine 25 mg PO 30 min before infusion, may repeat every 4 to 6 hours as needed for mild to moderate allergic reactions. Patient may decline.
- Methylprednisolone Sodium Succinate 40 mg IV push 20 minutes prior to infusion.
- Other: \_\_\_\_\_

### IV Flush Orders

- Peripheral: 0.9% Sodium Chloride 3 to 5 mL pre-/post-use.
- Implanted Port: 0.9% Sodium Chloride 5 to 10 mL pre-/post-use and 10 to 20 mL pre-/post-lab draw. Heparin (100 unit/mL) 3 to 5 mL post-use. For maintenance, heparin (100 unit/mL) 3 to 5 mL every 24 hr if accessed or weekly to monthly if not accessed.

### Lab Orders

- No labs ordered at this time.
- Other: \_\_\_\_\_

Skilled nurse to assess and administer and/or teach self-administration where appropriate via (IV/SubQ) access device as indicated above. Nurse will provide ongoing support as needed. Refill above ancillary orders as directed x 1 year.

If patient is seen within a provider led infusion clinic, Option Care Health's infusion reaction management policy, skilled nursing plan of treatment, and IV flush administration will be followed per provider oversight. No individual anaphylaxis kit will be dispensed.

*I certify that the use of the indicated treatment is medically necessary, and I will be supervising the patient's treatment.*

Prescriber Signature:

Date:

## Prescriber Information

Prescriber Name:	Phone:	Fax:
Address:	NPI:	
City, State:	Zip:	Office Contact:

Fax completed form, insurance information, and clinical documentation to: 713-983-4647

**CONFIDENTIAL HEALTH INFORMATION:** Healthcare information is personal information related to a person's healthcare. It is being faxed to you after appropriate authorization or under circumstances that do not require authorization. You are obligated to maintain it in a safe, secure, and confidential manner. Re-disclosure of this information is prohibited unless permitted by law or appropriate customer/patient authorization is obtained. Unauthorized re-disclosure or failure to maintain confidentiality could subject you to penalties described in federal and state laws. **IMPORTANT WARNING:** This message is intended for the use of the person or entity to whom it is addressed and may contain information that is privileged and confidential, the disclosure of which is governed by applicable law. If the reader of this message is not the intended recipient, or the employee or agent responsible for delivering it to the intended recipient, you are hereby notified that any dissemination, distribution or copying of this information is STRICTLY PROHIBITED. If you have received this message in error, please notify us immediately. Brand names are the property of their respective owners.

**Not valid for use for patients residing in Arizona, New York, and Wisconsin**