

OMVOH® (MIRIKIZUMAB) PRESCRIBER ORDER FORM

Patient Name:		Date of Birth:	Gender:	
Address:				
Phone:	Height:	<input type="checkbox"/> inches <input type="checkbox"/> cm	Weight:	<input type="checkbox"/> lbs <input type="checkbox"/> kg

Clinical Information

Primary Diagnosis Description:		ICD-10 Code:
TB Status:	<input type="checkbox"/> PPD (negative) – date:	<input type="checkbox"/> Active TB
	<input type="checkbox"/> Last Chest x-ray – date:	<input type="checkbox"/> Unknown
	<input type="checkbox"/> QuantiFERON or T Spot Assay result and date:	<input type="checkbox"/> Past positive TB infection, course taken:
Is this the first dose?	<input type="checkbox"/> Yes – date of first dose: _____	<input type="checkbox"/> No – date of last dose: _____

OmvoH® (Mirikizumab) Prescription

OmvoH® (Mirikizumab) refill as directed x 1 year

Ulcerative Colitis

Induction Dose: Infuse 300 mg IV over at least 30 minutes at Weeks 0, 4, and 8.

Maintenance Dose: Inject 200mg subcutaneously at week 12 and every 4 weeks thereafter (given as one injection of 200mg or two consecutive injections of 100mg).

Other: _____

Crohn's Disease

Induction Dose: Infuse 900 mg IV over at least 90 minutes at Weeks 0, 4, and 8.

Maintenance Dose: Inject 300mg subcutaneously at week 12 and every 4 weeks thereafter (given as two consecutive injections of 100mg and 200mg in any order).

Other: _____

After each infusion, the IV tubing will be flushed with 0.9% Sodium Chloride 30ml using a 50ml bag.

Ancillary Orders

Anaphylaxis Kit

Does this patient require an anaphylaxis kit?

Yes, with 1st dose Yes, with all doses

Dosage: Epinephrine 0.3 mg (> 30 kg), 0.15 mg (15 to 30 kg), or 0.01 mg/kg (< 15 kg) SUBQ or IM x 1; repeat x 1 in 5 to 15 min PRN.

Diphenhydramine 25 mg (> 30 kg) or 1.25 mg/kg (≤ 30 kg) IV or IM; repeat x 1 in 15 min PRN no improvement.

0.9% Sodium Chloride 500 mL (> 30 kg) or 250 mL (≤ 30 kg) IV at KVO rate PRN anaphylaxis.

Medication Orders

Other: _____

IV Flush Orders

Peripheral: 0.9% Sodium Chloride 3 to 5 mL pre-/post-use.

Implanted Port: 0.9% Sodium Chloride 5 to 10 mL pre-/post-use and 10 to 20 mL pre-/post-lab draw. Heparin (100 unit/mL) 3 to 5 mL post-use. For maintenance, heparin (100 unit/mL) 3 to 5 mL every 24 hr if accessed or weekly to monthly if not accessed.

Lab Orders

No labs ordered at this time.

Other: _____

Skilled nurse to administer doses intravenously in the home or alternate care setting. Refill above ancillary orders as directed x 1 year.

If patient is seen within a provider led infusion clinic, Option Care Health's infusion reaction management policy, skilled nursing plan of treatment, and IV flush administration will be followed per provider oversight. No individual anaphylaxis kit will be dispensed.

I certify that the use of the indicated treatment is medically necessary, and I will be supervising the patient's treatment.

Prescriber Signature:	Date:
-----------------------	-------

Prescriber Information

Prescriber Name:	Phone:	Fax:
Address:	NPI:	
City, State:	Zip:	Office Contact:

Fax completed form, insurance information, and clinical documentation to: 713-983-4647

CONFIDENTIAL HEALTH INFORMATION: Healthcare information is personal information related to a person's healthcare. It is being faxed to you after appropriate authorization or under circumstances that do not require authorization. You are obligated to maintain it in a safe, secure, and confidential manner. Re-disclosure of this information is prohibited unless permitted by law or appropriate customer/patient authorization is obtained. Unauthorized re-disclosure or failure to maintain confidentiality could subject you to penalties described in federal and state laws. **IMPORTANT WARNING:** This message is intended for the use of the person or entity to whom it is addressed and may contain information that is privileged and confidential, the disclosure of which is governed by applicable law. If the reader of this message is not the intended recipient, or the employee or agent responsible for delivering it to the intended recipient, you are hereby notified that any dissemination, distribution or copying of this information is STRICTLY PROHIBITED. If you have received this message in error, please notify us immediately. Brand names are the property of their respective owners.

Not valid for use for patients residing in Arizona, New York, and Wisconsin