POTELIGEO® (mogamulizumab-kpkc) PRESCRIBER ORDER FORM							
Patient Name:				Date of Birth:			
Address:							
Phone:			Height:	☐ inches ☐ cn	Weight:	☐ lbs. ☐ kg	
Clinical Information							
Primary Diagnosis Description: ICD-10 Code:							
Date of first dose received (Day 1):							
POTELIGEO® (mogamulizumab-kpkc) Prescription POTELIGEO® (mogamulizumab-kpkc) Refill as directed x1 year. First dose to be administered in prescriber's office.							
1 mg/kg administered as an intravenous infusion over at least 60 minutes. Administer on days 8, 15, and 22 of the first 28-day cycle, then on days 1 and 15 of each subsequent 28-day cycle until disease progression or unacceptable toxicity. Administer within 2 days of the scheduled dose.							
Calculate the dose (mg/kg) and number of single dose vials needed to prepare the infusion solution based on patient weight. Whole vials will be dispensed based on calculated dose. Discard any unused portion left in the vial.							
Ancillary Orders Anaphylaxis Orders							
 Epinephrine 0.3 mg (> 30 kg), 0.15 mg (15 to 30 kg), or 0.01 mg/kg (< 15 kg) SubQ or IM x 1; repeat x 1 in 5 to 15 min PRN. Diphenhydramine 25 mg (> 30 kg) or 1.25 mg/kg (≤ 30 kg) IV or IM; repeat x 1 in 15 min PRN no improvement. 0.9% Sodium Chloride 500 mL (> 30 kg) or 250 mL (≤ 30 kg) IV at KVO rate PRN anaphylaxis. 							
Pre-Medication Orders							
☐ Acetaminophen 650 mg PO 30 min before infusion.							
☐ Diphenhydramine 25 mg PO 30 min before infusion.							
□ Other:							
IV Flush Orders							
	<u>Peripheral:</u>	0.9% Sodium Chloride	e 2 to 3 mL pre-	/post-use.			
	PICC and Central Tunneled/Non- Tunneled:	0.9% Sodium Chloride 5 to 10 pre-/post-use, 5 mL pre-lab draw and 10 ml post-lab draw. Heparin (10 unit/mL) 5 mL or (100 unit/mL post-use. For maintenance, Heparin (10 unit/mL) 5 mL or (100 unit/mL)3 mL every 24 hr.					
	Implanted Port:	0.9% Sodium Chloride 5 to 10 mL pre-/post-use and 10 to 20 mL pre-/post-lab draw. Heparin (100 unit/mL) 3 to 5 mL post-use. For maintenance, heparin (100 unit/mL) 3 to 5 mL every 24 hr. if accessed or weekly to monthly if not accessed.					
	Valved Catheters:	0.9% Sodium Chloride 5 to 10 mL pre-/post-use and 10 to 20 mL pre-/post-lab draw. For maintenance, 0.9% Sodium Chloride NS 5 to 10 ml at least weekly					
Lab Orders							
	No labs ordered at this time.						
Other:Skilled nurse to assess and administer via access device as indicated above. Nurse will provide ongoing support as needed.							
Refill above ancillary orders as directed x 1 year.							
I certify that the use of the indicated treatment is medically necessary, and I will be supervising the patient's treatment.							
Prescriber Signature:					Date:		
		criber Informat	ion				
Prescriber Name	:		Phone:		Fax:		
Address:			NPI:				
City, State: Zip: Office Contact:							

CONFIDENTIAL HEALTH INFORMATION: Healthcare information is personal information related to a person's healthcare. It is being faxed to you after appropriate authorization or under circumstances that do not require authorization. You are obligated to maintain it in a safe, secure, and confidential manner. Re-disclosure of this information is prohibited unless permitted by law or appropriate customer/patient authorization is obtained. Unauthorized re-disclosure or failure to maintain confidentiality could subject vous to penalties described in federal and state laws. IMPORTANT WARNING: This message is intended for the use of the person or entity to whom it is addressed and may contain information that is privileged and confidential, the disclosure of which is governed by applicable law. If the reader of this message is not the intended recipient, or the employee or agent responsible for delivering it to the intended recipient, you are hereby notified that any dissemination, distribution or copying of this information is STRICTLY PROHIBITED. If you have received this message in error, please notify us immediately. Brand names are the property of their respective owners.