


TOCILIZUMAB (ACTEMRA®) PRESCRIBER ORDER FORM

Fax completed form, insurance information, and clinical documentation to:

	Patient Name:		Date of Birth:	
	Address:			
	Patient Phone:	Height:	<input type="checkbox"/> inches <input type="checkbox"/> cm	Weight:

Clinical Information

Primary Diagnosis Description:		ICD-10 Code:	
Is this the first dose? <input type="checkbox"/> Yes – date of first dose: <input type="checkbox"/> No – date of next dose due:		Hepatitis B Status:	Titer Date:
TB Status: <input type="checkbox"/> PPD (negative) – date: <input type="checkbox"/> Last chest x-ray – date: <input type="checkbox"/> Past positive TB infection, course taken:		<input type="checkbox"/> Active TB <input type="checkbox"/> Unknown	<input type="checkbox"/> Positive <input type="checkbox"/> Negative

Tocilizumab (Actemra®) Prescription**Tocilizumab (Actemra®) refill as directed x 1 year**

- Infuse ____ mg/kg IV over 60 minutes every 4 weeks – max 800 mg.
 Inject 162 mg once every week or every other week.
 Other: _____

Ancillary Orders**Anaphylaxis Kit**

- Dosage:**
- Epinephrine 0.3 mg (> 30 kg), 0.15 mg (15 to 30 kg), or 0.01 mg/kg (< 15 kg) SQ or IM x 1; repeat x 1 in 5 to 15 min PRN.
 - Diphenhydramine 25 mg (> 30 kg) or 1.25 mg/kg (≤ 30 kg) IV or IM; repeat x 1 in 15 min PRN no improvement.
 - Normal saline 500 mL (> 30 kg) or 250 mL (≤ 30 kg) IV at KVO rate PRN anaphylaxis. Patients ≤ 30 kg, infuse over 2 to 4 hours PRN headache rated > 5 on pain scale.

Medication Orders

- Acetaminophen 650 mg PO 30 min before infusion, may repeat every 3 to 4 hours as needed for fever or mild discomfort. Patient may use own supply or patient may decline.
- Diphenhydramine 25 mg PO 30 min before infusion, may repeat every 4 to 6 hours as needed for mild to moderate allergic reactions. Patient may use own supply or patient may decline.
- Methylprednisolone 40 mg IV push 20 minutes prior to infusion.
- Other: _____

IV Flush Orders

- Peripheral: NS 2 to 3 mL pre-/post-use.
- Implanted Port: NS 5 to 10 mL pre-/post-use and 10 to 20 mL pre-/post-lab draw. Heparin (100 unit/mL) 3 to 5 mL post-use. For maintenance, heparin (100 unit/mL) 3 to 5 mL every 24 hr if accessed or weekly to monthly if not accessed.

Lab Orders

- No labs ordered at this time.
- Other: _____

Skilled nurse to administer doses intravenously in the home or alternate care setting. Refill above ancillary orders as directed x 1 year.

I certify that the use of the indicated treatment is medically necessary and I will be supervising the patient's treatment.

Prescriber Signature:**Date:****Prescriber Information**

Prescriber Name:	Phone:	Fax:
Address:	NPI:	
City, State:	Zip:	Office Contact:

CONFIDENTIAL HEALTH INFORMATION: Healthcare information is personal information related to a person's healthcare. It is being faxed to you after appropriate authorization or under circumstances that don't require authorization. You are obligated to maintain it in a safe, secure, and confidential manner. Re-disclosure of this information is prohibited unless permitted by law or appropriate customer/patient authorization is obtained. Unauthorized re-disclosure or failure to maintain confidentiality could subject you to penalties described in federal and state laws. **IMPORTANT WARNING:** This message is intended for the use of the person or entity to whom it is addressed and may contain information that is privileged and confidential, the disclosure of which is governed by applicable law. If the reader of this message is not the intended recipient, or the employee or agent responsible for delivering it to the intended recipient, you are hereby notified that any dissemination, distribution or copying of this information is STRICTLY PROHIBITED. If you have received this message in error, please notify us immediately. Brand names are the property of their respective owners.