VEDOLIZUM	IAB (ENTYVIO®) PRESCRIBER	ORDER FORI	M					
Patient Name:			Date of Birth:			Gender:	Gender:	
Address:			<u>'</u>				<u> </u>	
Phone:		н	leight:	□ inc	hes $\square$ cm	Weigl	ht:	☐ lbs ☐ kg
		Clinica	al Informatio	on				
Primary Diagno	sis Description:				IC	D-10 Co	de:	
Is this the first o	<b>dose?</b>			□ NO – Da	te of last dos	e:		
		Vedolizumab (						
Vedolizumab (Entyvio®) refill as directed x 1 year. Please select <u>one</u> of the options below:								
	☐ Option 1:	☐ Option 2:			☐ Option 3	3:		
Induction	Infuse 300 mg IV over at least 30	Infuse Entyvio	_				ng <b>IV</b> over at	least
Maintananaa	minutes on Weeks <b>0, 2, 6</b>	30 minutes at						un als C
Maintenance	Infuse 300 mg <b>IV</b> over at least 30 minutes every 8 weeks thereafter	Inject Entyvio 14, then every	-		then every			eek 6,
	·	<u> </u>	,		<u> </u>			
Other:								
	ntity sufficient of Entyvio® 300mg vials will be flushed with NS 30ml using a 50ml bag.		he patient to	complete the p	rescribed treat	ment pla	ın. In addition,	after each infusion
	quantity sufficient of Entyvio® 108mg Prefi		ispensed to fu	Ifill prescribed	treatment plan	ո.		
		Anci	illary Orders					
Anaphylaxis K								
	infusion dose, would you like Option  ☐ Yes ☐ No	Care Health to p	orovide an ar	naphylaxis kit	with the 1 <sup>st</sup> d	lose?		
•	Epinephrine 0.3 mg (> 30 kg), 0.15 mg Diphenhydramine 25 mg (> 30 kg) or	1.25 mg/kg (≤ 3	0 kg – 25mg	max dose) IV	or IM; repea			
	0.9% Sodium Chloride 500 mL (> 30 kg to establish peripheral IV access as				anaphylaxis.			
Medication O		necucu to mana	ige anapriyia	AIJ.				
	etaminophen 650 mg PO 30 min before infusion, may repeat every 3 to 4 hours as needed for fever or mild discomfort. Patient by decline.							
	henhydramine 25 mg PO 30 min before infusion, may repeat every 4 to 6 hours as needed for mild to moderate allergic reactions. ient may decline.							
□ Me	thylprednisolone Sodium Succinate 40 mg IV push 20 minutes prior to infusion.							
□ Ot	ther:							
IV Flush Orde								
	eripheral: 0.9% Sodium Chloride 2 to 3 mL pre-/post-use.  nplanted Port: 0.9% Sodium Chloride 5 to 10 mL pre-/post-use and 10 to 20 mL pre-/post-lab draw. Heparin (100 unit/mL) to 5 mL post-useFor maintenance, heparin (100 unit/mL) 3 to 5 mL every 24 hr if accessed or weekly to monthly if not accessed.							
Lab Orders	a labo and anad at this tim							
_	o labs ordered at this time. ther:							
<del></del>	o assess and administer and/or teach	self-administrat	ion where a	ppropriate via	a (IV/SubQ) a	ccess de	vice as indica	ated above. Nurse
•	ngoing support as needed. Refill above	•		•	•			
	en within a provider led infusion clinion will be followed per provider oversig				_	policy, s	killed nursing	g plan of treatment,
	certify that the use of the indicated tre					the pat	tient's treatm	nent.
	•		,					
Prescriber Signa	ature:	Prescrib	oer Informat	ion		υ 	ate:	
Prescriber Name:			Phone:	- IOII	Fax:			
Address:			NPI:		<u>                                  </u>			
City, State: Zip:			Office Contact:					
Fax completed	form, insurance information, and clin	ical documentat	tion to: 713	3-983-4647				
	HINFORMATION: Healthcare information is personal infobligated to maintain it in a safe, secure, and confidentia							

Unauthorized re-disclosure or failure to maintain this a sare, secure, and confidentiality could subject you to penalties described in federal and state laws. IMPORTANT WARNING: This message is intended for the use of the person or entity to whom it is addressed and may contain information that is privileged and confidential, the disclosure of which is governed by applicable law. If the reader of this message is not the intended recipient, or the employee or agent responsible for delivering it to the intended recipient, you are hereby notified that any dissemination, distribution or copying of this information is STRICTLY PROHIBITED. If you have received this message in error, please notify us immediately. Brand names are the property of their respective owners.